


# Nattokinase NSK-SD®

*The original Superior Oral fibrinolytic enzyme  
For Healthy Circulation\**



- **Enhances fibrinolytic activity\***
- **Supports the body's regulation of blood viscosity\***
- **Supports healthy coagulation of blood within normal levels\***

## ***As the Leading Doctor's Nattokinase, NSK-SD® Sets the Standard!***

**Nattokinase** is an enzyme derived from boiled soybeans and *Bacillus subtilis natto*. Research has shown nattokinase to support healthy coagulation of blood within normal levels and enhance fibrinolytic activity.\* Nattokinase NSK-SD® enhances the body's natural fibrinolytic activity, the dissolution of the essential portion of the blood clot or thrombus.\* Maintenance of optimal functioning of the body's fibrinolytic/thrombolytic mechanisms may benefit the function of the cardiovascular system and the brain.\*

**Allergy Research Group's NSK-SD®** was the first Nattokinase in the U.S. market, and it has

established standardization and quality levels for all nattokinase, with comprehensive safety studies and proven potency. It is vegetarian, non-irradiated, and free of vitamin K2. NSK-SD® has two Japanese and three U.S. patents, and is recognized by the JHFA (Japan Health and Nutrition Food Authorization) and JNKA (Japan Nattokinase Association) as authentic nattokinase. NSK-SD® is produced from non-genetically modified soybeans using a selected, patented strain of *Bacillus subtilis natto*. Each batch is tested to ensure potency.

**Allergy Research Group®**

**Phone: 800-545-9960/510-263-2000 • Fax: 800-688-7426/510-263-2100**

**[www.allergyresearchgroup.com](http://www.allergyresearchgroup.com)**



NSK-SD® is a trademark of Japan BioScience Laboratory.

\*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

# Nattokinase NSK-SD<sup>®</sup>



## **Nattokinase 100 mg NSK-SD<sup>™</sup>, #75370, 75380**

Each 1 softgel contains:

Nattokinase NSK-SD<sup>™</sup> (2000 FU) . . . 100 mg

**Other ingredients:** Soybean oil, gelatin, glycerin, soybean lecithin, glycerin fatty acid ester, beeswax.

**Suggested use:** As a dietary supplement, 1 softgel two times daily initial dose, or as directed by a healthcare practitioner. May be taken with or without food. Take with 8-10 ounces of water. If taken with anticoagulant drugs, use under medical supervision. Contraindicated in any condition associated with bleeding.



## **Nattokinase 50 mg NSK-SD<sup>™</sup>, #75281, 75291**

Each 2 vegetarian capsules contain:

Nattokinase NSK-SD<sup>™</sup> (2000 FU) . . . 100 mg

**Other ingredients:** Hydroxypropyl methylcellulose, cellulose, L-leucine.

**Suggested use:** As a dietary supplement, 2 capsules two times daily initial dose, or as directed by a healthcare practitioner. May be taken with or without food. Take with 8-10 ounces of water. If taken with anticoagulant drugs, use under medical supervision. Contraindicated in any condition associated with bleeding.



## **Nattokinase 36 mg NSK-SD<sup>™</sup>, #74750, 75100**

Each 2 softgels contain:

Nattokinase NSK-SD<sup>™</sup> (1440 FU) . . . 72 mg

**Other ingredients:** Soybean oil, gelatin, glycerin, soybean lecithin, glycerin fatty acid ester, beeswax.

**Suggested use:** As a dietary supplement, 1 or 2 softgels three times daily, or as directed by a healthcare practitioner. May be taken with or without food. Take with 8-10 ounces of water. If taken with anticoagulant drugs, use under medical supervision. Contraindicated in any condition associated with bleeding.

**Allergy Research Group<sup>®</sup>**

**Phone: 800-545-9960/510-263-2000 • Fax: 800-688-7426/510-263-2100**

**[www.allergyresearchgroup.com](http://www.allergyresearchgroup.com)**

